

# Lecture 4

## Orthonormal sets of vectors and $QR$ factorization

- orthonormal sets of vectors
- Gram-Schmidt procedure,  $QR$  factorization
- orthogonal decomposition induced by a matrix

# Orthonormal set of vectors

set of vectors  $u_1, \dots, u_k \in \mathbf{R}^n$  is

- *normalized* if  $\|u_i\| = 1, i = 1, \dots, k$   
( $u_i$  are called *unit vectors* or *direction vectors*)
- *orthogonal* if  $u_i \perp u_j$  for  $i \neq j$
- *orthonormal* if both

**slang:** we say ' $u_1, \dots, u_k$  are orthonormal vectors' but orthonormality (like independence) is a property of a set of vectors, not vectors individually

in terms of  $U = [u_1 \ \cdots \ u_k]$ , orthonormal means

$$U^T U = I_k$$

- orthonormal vectors are independent  
(multiply  $\alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_k u_k = 0$  by  $u_i^T$ )
- hence  $u_1, \dots, u_k$  is an orthonormal basis for

$$\text{span}(u_1, \dots, u_k) = \mathcal{R}(U)$$

- **warning:** if  $k < n$  then  $UU^T \neq I$  (since its rank is at most  $k$ )  
(more on this matrix later . . . )

# Geometric properties

suppose columns of  $U = [u_1 \ \cdots \ u_k]$  are orthonormal

if  $w = Uz$ , then  $\|w\| = \|z\|$

- multiplication by  $U$  does not change norm
- mapping  $w = Uz$  is *isometric*: it preserves distances
- simple derivation using matrices:

$$\|w\|^2 = \|Uz\|^2 = (Uz)^T(Uz) = z^T U^T U z = z^T z = \|z\|^2$$

- *inner products* are also preserved:  $\langle Uz, U\tilde{z} \rangle = \langle z, \tilde{z} \rangle$
- if  $w = Uz$  and  $\tilde{w} = U\tilde{z}$  then

$$\langle w, \tilde{w} \rangle = \langle Uz, U\tilde{z} \rangle = (Uz)^T (U\tilde{z}) = z^T U^T U \tilde{z} = \langle z, \tilde{z} \rangle$$

- norms and inner products preserved, so *angles* are preserved:  
 $\angle(Uz, U\tilde{z}) = \angle(z, \tilde{z})$
- thus, multiplication by  $U$  preserves inner products, angles, and distances

## Orthonormal basis for $\mathbf{R}^n$

- suppose  $u_1, \dots, u_n$  is an orthonormal *basis* for  $\mathbf{R}^n$
- then  $U = [u_1 \cdots u_n]$  is called **orthogonal**: it is square and satisfies  $U^T U = I$

(you'd think such matrices would be called *orthonormal*, not *orthogonal*)

- it follows that  $U^{-1} = U^T$ , and hence also  $U U^T = I$ , *i.e.*,

$$\sum_{i=1}^n u_i u_i^T = I$$

# Expansion in orthonormal basis

suppose  $U$  is orthogonal, so  $x = UU^T x$ , *i.e.*,

$$x = \sum_{i=1}^n (u_i^T x) u_i$$

- $u_i^T x$  is called the *component* of  $x$  in the direction  $u_i$
- $a = U^T x$  *resolves*  $x$  into the vector of its  $u_i$  components
- $x = Ua$  *reconstitutes*  $x$  from its  $u_i$  components

- $x = Ua = \sum_{i=1}^n a_i u_i$  is called the  $(u_i)$  *expansion* of  $x$

the identity  $I = UU^T = \sum_{i=1}^n u_i u_i^T$  is sometimes written (in physics) as

$$I = \sum_{i=1}^n |u_i\rangle\langle u_i|$$

since

$$x = \sum_{i=1}^n |u_i\rangle\langle u_i|x\rangle$$

(but we won't use this notation)



# Geometric interpretation

if  $U$  is orthogonal, then transformation  $w = Uz$

- preserves *norm* of vectors, *i.e.*,  $\|Uz\| = \|z\|$
- preserves *angles* between vectors, *i.e.*,  $\angle(Uz, U\tilde{z}) = \angle(z, \tilde{z})$

examples:

- rotations (about some axis)
- reflections (through some plane)

**Example:** rotation by  $\theta$  in  $\mathbf{R}^2$  is given by



$$y = U_\theta x, \quad U_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

since  $e_1 \rightarrow (\cos \theta, \sin \theta)$ ,  $e_2 \rightarrow (-\sin \theta, \cos \theta)$

reflection across line  $x_2 = x_1 \tan(\theta/2)$  is given by

$$y = R_\theta x, \quad R_\theta = \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix}$$

since  $e_1 \rightarrow (\cos \theta, \sin \theta)$ ,  $e_2 \rightarrow (\sin \theta, -\cos \theta)$



can check that  $U_\theta$  and  $R_\theta$  are orthogonal

# Gram-Schmidt procedure


- given independent vectors  $a_1, \dots, a_k \in \mathbf{R}^n$ , G-S procedure finds orthonormal vectors  $q_1, \dots, q_k$  s.t.

$$\text{span}(a_1, \dots, a_r) = \text{span}(q_1, \dots, q_r) \quad \text{for } r \leq k$$

- thus,  $q_1, \dots, q_r$  is an orthonormal basis for  $\text{span}(a_1, \dots, a_r)$
- rough idea of method: first *orthogonalize* each vector w.r.t. previous ones; then *normalize* result to have norm one

# Gram-Schmidt procedure

- step 1a.  $\tilde{q}_1 := a_1$
- step 1b.  $q_1 := \tilde{q}_1 / \|\tilde{q}_1\|$  (normalize)
- step 2a.  $\tilde{q}_2 := a_2 - (q_1^T a_2)q_1$  (remove  $q_1$  component from  $a_2$ )
- step 2b.  $q_2 := \tilde{q}_2 / \|\tilde{q}_2\|$  (normalize)
- step 3a.  $\tilde{q}_3 := a_3 - (q_1^T a_3)q_1 - (q_2^T a_3)q_2$  (remove  $q_1, q_2$  components)
- step 3b.  $q_3 := \tilde{q}_3 / \|\tilde{q}_3\|$  (normalize)
- etc.



for  $i = 1, 2, \dots, k$  we have

$$\begin{aligned}
 a_i &= (q_1^T a_i)q_1 + (q_2^T a_i)q_2 + \cdots + (q_{i-1}^T a_i)q_{i-1} + \|\tilde{q}_i\|q_i \\
 &= r_{1i}q_1 + r_{2i}q_2 + \cdots + r_{ii}q_i
 \end{aligned}$$

(note that the  $r_{ij}$ 's come right out of the G-S procedure, and  $r_{ii} \neq 0$ )

# QR decomposition

written in matrix form:  $A = QR$ , where  $A \in \mathbf{R}^{n \times k}$ ,  $Q \in \mathbf{R}^{n \times k}$ ,  $R \in \mathbf{R}^{k \times k}$ :

$$\underbrace{\begin{bmatrix} a_1 & a_2 & \cdots & a_k \end{bmatrix}}_A = \underbrace{\begin{bmatrix} q_1 & q_2 & \cdots & q_k \end{bmatrix}}_Q \underbrace{\begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1k} \\ 0 & r_{22} & \cdots & r_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{kk} \end{bmatrix}}_R$$

- $Q^T Q = I_k$ , and  $R$  is upper triangular & invertible
- called **QR decomposition** (or factorization) of  $A$
- usually computed using a variation on Gram-Schmidt procedure which is less sensitive to numerical (rounding) errors
- columns of  $Q$  are orthonormal basis for  $\mathcal{R}(A)$

# General Gram-Schmidt procedure

- in basic G-S we assume  $a_1, \dots, a_k \in \mathbf{R}^n$  are independent
- if  $a_1, \dots, a_k$  are dependent, we find  $\tilde{q}_j = 0$  for some  $j$ , which means  $a_j$  is linearly dependent on  $a_1, \dots, a_{j-1}$
- modified algorithm: when we encounter  $\tilde{q}_j = 0$ , skip to next vector  $a_{j+1}$  and continue:


```
 $r = 0;$   
for  $i = 1, \dots, k$   
{  
     $\tilde{a} = a_i - \sum_{j=1}^r q_j q_j^T a_i;$   
    if  $\tilde{a} \neq 0$  {  $r = r + 1; q_r = \tilde{a} / \|\tilde{a}\|;$  }  
}
```



on exit,

- $q_1, \dots, q_r$  is an orthonormal basis for  $\mathcal{R}(A)$  (hence  $r = \mathbf{Rank}(A)$ )
- each  $a_i$  is linear combination of previously generated  $q_j$ 's

in matrix notation we have  $A = QR$  with  $Q^T Q = I_r$  and  $R \in \mathbf{R}^{r \times k}$  in *upper staircase form*:



'corner' entries (shown as  $\times$ ) are nonzero

can permute columns with  $\times$  to front of matrix:

$$A = Q[\tilde{R} \ S]P$$

where:

- $Q^T Q = I_r$
- $\tilde{R} \in \mathbf{R}^{r \times r}$  is upper triangular and invertible
- $P \in \mathbf{R}^{k \times k}$  is a permutation matrix  
(which moves forward the columns of  $a$  which generated a new  $q$ )

# Applications

- directly yields orthonormal basis for  $\mathcal{R}(A)$
- yields factorization  $A = BC$  with  $B \in \mathbf{R}^{n \times r}$ ,  $C \in \mathbf{R}^{r \times k}$ ,  $r = \mathbf{Rank}(A)$
- to check if  $b \in \text{span}(a_1, \dots, a_k)$ : apply Gram-Schmidt to  $[a_1 \ \cdots \ a_k \ b]$
- staircase pattern in  $R$  shows which columns of  $A$  are dependent on previous ones

works incrementally: one G-S procedure yields  $QR$  factorizations of  $[a_1 \ \cdots \ a_p]$  for  $p = 1, \dots, k$ :

$$[a_1 \ \cdots \ a_p] = [q_1 \ \cdots \ q_s]R_p$$

where  $s = \mathbf{Rank}([a_1 \ \cdots \ a_p])$  and  $R_p$  is leading  $s \times p$  submatrix of  $R$

## 'Full' QR factorization

with  $A = Q_1 R_1$  the QR factorization as above, write

$$A = [ Q_1 \quad Q_2 ] \begin{bmatrix} R_1 \\ 0 \end{bmatrix}$$

where  $[Q_1 \quad Q_2]$  is orthogonal, *i.e.*, columns of  $Q_2 \in \mathbf{R}^{n \times (n-r)}$  are orthonormal, orthogonal to  $Q_1$

to find  $Q_2$ :

- find any matrix  $\tilde{A}$  s.t.  $[A \quad \tilde{A}]$  is full rank (*e.g.*,  $\tilde{A} = I$ )
- apply general Gram-Schmidt to  $[A \quad \tilde{A}]$
- $Q_1$  are orthonormal vectors obtained from columns of  $A$
- $Q_2$  are orthonormal vectors obtained from extra columns ( $\tilde{A}$ )

*i.e.*, any set of orthonormal vectors can be *extended* to an orthonormal basis for  $\mathbf{R}^n$

$\mathcal{R}(Q_1)$  and  $\mathcal{R}(Q_2)$  are called *complementary subspaces* since

- they are orthogonal (*i.e.*, every vector in the first subspace is orthogonal to every vector in the second subspace)
- their sum is  $\mathbf{R}^n$  (*i.e.*, every vector in  $\mathbf{R}^n$  can be expressed as a sum of two vectors, one from each subspace)

this is written

- $\mathcal{R}(Q_1) \overset{\perp}{+} \mathcal{R}(Q_2) = \mathbf{R}^n$
- $\mathcal{R}(Q_2) = \mathcal{R}(Q_1)^\perp$  (and  $\mathcal{R}(Q_1) = \mathcal{R}(Q_2)^\perp$ )  
(each subspace is the *orthogonal complement* of the other)

we know  $\mathcal{R}(Q_1) = \mathcal{R}(A)$ ; but what is its orthogonal complement  $\mathcal{R}(Q_2)$ ?

## Orthogonal decomposition induced by $A$

from  $A^T = \begin{bmatrix} R_1^T & 0 \end{bmatrix} \begin{bmatrix} Q_1^T \\ Q_2^T \end{bmatrix}$  we see that

$$A^T z = 0 \iff Q_1^T z = 0 \iff z \in \mathcal{R}(Q_2)$$

so  $\mathcal{R}(Q_2) = \mathcal{N}(A^T)$

(in fact the columns of  $Q_2$  are an orthonormal basis for  $\mathcal{N}(A^T)$ )

we conclude:  $\mathcal{R}(A)$  and  $\mathcal{N}(A^T)$  are *complementary subspaces*:

- $\mathcal{R}(A) \overset{\perp}{+} \mathcal{N}(A^T) = \mathbf{R}^n$  (recall  $A \in \mathbf{R}^{n \times k}$ )
- $\mathcal{R}(A)^\perp = \mathcal{N}(A^T)$  (and  $\mathcal{N}(A^T)^\perp = \mathcal{R}(A)$ )
- called *orthogonal decomposition (of  $\mathbf{R}^n$ ) induced by  $A \in \mathbf{R}^{n \times k}$*

- every  $y \in \mathbf{R}^n$  can be written uniquely as  $y = z + w$ , with  $z \in \mathcal{R}(A)$ ,  $w \in \mathcal{N}(A^T)$  (we'll soon see what the vector  $z$  is . . . )
- can now prove most of the assertions from the linear algebra review lecture
- switching  $A \in \mathbf{R}^{n \times k}$  to  $A^T \in \mathbf{R}^{k \times n}$  gives decomposition of  $\mathbf{R}^k$ :

$$\mathcal{N}(A) \overset{\perp}{+} \mathcal{R}(A^T) = \mathbf{R}^k$$